

Stephen Biles

186 CR 101

Port Lavaca, TX 77979

# PEST MANAGEMENT NEWS

Calhoun, Refugio & Victoria Counties

(o) 361-552-3324

(m) 361-920-1138

biles-sp@tamu.edu

VOLUME 10

ISSUE 13

July 24, 2014

## Grain Sorghum

Grain fields continue to be harvested and yields have vary as widely as the rainfall from less than 3000 lbs/A to more than 7000 lbs/A. Most of the sorghum is at hard dough or black layer and beyond the damage window for panicle feeding stink bugs and headworms. A few young fields are still blooming and remain susceptible to sorghum midge, stink bugs and headworms.

All sorghum fields are susceptible to sugarcane aphids. Some sorghum producing areas have begun to call this the “white” sugarcane aphid to prevent confusion with the yellow sugarcane aphid. If you read articles about white sugarcane aphids in Louisiana, Arkansas, Mississippi or other states, they are the same aphid we are dealing with.

Harvesting problems occurring in parts of Refugio County caused by the sugarcane aphid and the honeydew it produces. The fields had sugarcane aphids on the flag leaf at the time of glyphosate applications and when the leaf died, the aphids moved to the head. In some fields, 30-40% of sorghum heads were infested with aphids.

We saw aphids move to the head in an insecticide trial near Port Lavaca, applied the same day as glyphosate was applied to the field, including the whole test area. By 10 days after treatment, some treatments averaged 60+ aphids in the head. These were producing sufficient honeydew to make the grain and upper leaves sticky with honey dew potentially causing harvest problems. Fortunately, this infestation was only in the field margins and not throughout the field.

A second trial is currently underway near Victoria to evaluate the combined and separate use of Transform Insecticide and RoundUp Powermax. More details will be available next week as we wait to see if the surviving aphids move to the head.

Honeydew from the aphids has also been observed reducing the efficacy of glyphosate applications causing some recommendations of increased rates of application. I have heard some recommendations of using sodium chlorate instead of glyphosate as a harvest aid. This product brings a lot of issues with regard to corrosion of equipment but may be a viable alternative if the product can be found.



Sugarcane Aphids on  
Mature Sorghum



Honeydew Shine on  
Sorghum Flag Leaf

## Cotton

Most cotton fields are past the damage window for stink bugs and Verde plant bugs (cutout (<5 NAWF) plus 350 Heat Units). We have some fields nearing the time to apply harvest aids.

When deciding on defoliation, look at heat units from cutout, percent open, and nodes above cracked boll. Proper timing of defoliation can be accomplished at cutout plus 850 heat units, >60% open bolls and/or 4 nodes above cracked boll to the highest harvestable boll.

We have found some worm damage in cotton bolls of Bt cotton. This is not uncommon, as Bollgard II, Widestrike and TwinLink cotton will control 90-95% of cotton bollworms. Usually this provides sufficient control of the worm population. In our counts, we have found less than 1% of bolls fed upon by bollworms.



Cotton Bollworm  
Feeding Damage

## Soybeans

Soybean fields are also closing in on harvest. I noticed James Grichar hauling his plot combine to his Port Lavaca variety trial. Stink bugs are still being found in soybean fields. Continue to monitor for stink bugs until bean maturity.

## Support for the 2014 IPM Program comes from the following:

Woodsboro Farmer's Cooperative  
Moreman Coop  
Hlavinka Equipment  
Numerous Producers

South Texas Cotton and Grain Association  
Helena Chemical  
Welfab

## Field Research is sponsored by:

- Cotton Incorporated
- AMVAC
- Syngenta
- Bayer CropScience
- Dow AgroScience
- Americot

To receive this newsletter via Text Message, text "Follow @Midcoastipm" to the number **40404**

To receive this newsletter via email, contact me at [biles-sp@tamu.edu](mailto:biles-sp@tamu.edu).

Also available at

- Twitter: @midcoastipm
- Facebook: Mid-Coast IPM

<http://calhoun.agrilife.org/newsletters/ipm-newsletter/>

[http://www.tpma.org/newsletters/coastal\\_middle/TOC.htm](http://www.tpma.org/newsletters/coastal_middle/TOC.htm)