

July 12, 2018, No. 18-5

Bill Ree, Extension Program Specialist III – IPM
AgriLife Extension South Campus
1470 William D. Fitch Parkway, 2150 TAMU
College Station, TX 77843-2150
979-458-0335 – office
w-ree@tamu.edu

TEXAS PECAN GROWERS ASSOCIATION
4348 Carter Creek, Suite 101 Bryan, TX 77802
Ph: 979-846-3285; Fax: 979-846-1752
pecans@tpga.org , www.tpga.org

TPGA CONFERENCE

The TPGA conference is just around the corner (July 15 – 18) and once again, TPGA has put together a great conference agenda and tour so I hope you will be attending:
https://www.tpga.org/upload/text/TPGA_Conference_Agenda:Educational_Program.pdf

During the conference I will have a booth near the registration desk where you will be able to find me during the breaks. Hope to see you next week.

INSECTS

Most of the pest issues I am seeing and/or receiving reports on are foliage pests, primarily black pecan aphid and some mite issues. I've always been told that there are two crops on a tree, the nut crop and the foliage which represents the next years crop. In order to have a good crop you need to protect the foliage and black aphids and mites can be a serious problem.

Black pecan aphid: Watch for black aphids in crowded areas of the orchard and in the interior of the canopy. I have also seen higher infestations on Pawnee.




Figure 1. Black pecan aphids

Insecticide recommendations for black aphids can be found in a table at the end of the newsletter. The treatment threshold is an average of 3 per compound leaf. This aphid can be found on both the upper and lower surfaces of the leaf and damage can be characterized by the rectangular yellow areas that stop abruptly at a secondary leaf vein as shown in Figure 1.

Pecan leaf scorch mites: As the growing season is now entering the hot dry months of mid-summer, one important foliage pest that is showing up is the pecan leaf scorch mite, *Eotetranychus hicoriae* (McGregor). Although

there are numerous species (27 species covering eight families of phytophagous and predatory species) associated with pecan, the pecan leaf scorch mite, is one of if not the most important.

Scorch mites are present all year and can be held below damaging population levels by predators and environmental conditions, however, pesticide use and hot dry conditions can lead to a population build up which will require management. As with all foliage pests, early recognition of a problem along with management actions is the key to preventing significant damage.

Scorch mites have multiple generations per year and development time from egg to adult is very short, 5 to 15 days under favorable conditions so damaging populations can arise very quickly. Feeding by scorch mites will result in brownish or liver colored areas along the mid-vein of leaflets as shown in the attached picture. This scorching should not be confused with leaf scorch along leaflet margins caused by nutrient imbalance or pecan bacterial leaf scorch.

Damaging infestations generally start in the lower and interior areas of the canopy then spread outward as populations increase. If a damaging infestation is detected, then surrounding trees should be checked. If an infestation is localized then a spot treatment might be applied.

When management is needed, there are numerous products to choose from. Products for pecan leaf scorch mites can be found in the table at the end of this letter. As with any pesticide, always read and follow directions for application, restrictions and safety.



Figure 2. Pecan leaf scorch mite damage

Walnut caterpillar: Although I have not received any reports lately, second generation walnut caterpillar is active at this time. As I mentioned in my previous letter, recognition of the early stages of an infestation is critical to preventing a major loss. For Texas there can be 2 to 3 generations per year with 245 frost free days being the dividing line. Usually trees damaged by one generation will not be infested by the next generation because the foliage will not be at the right maturity.



Figure 3. 5th instar walnut caterpillar

**2018 COUNTY/ STATE/REGIONAL
MEETINGS/EVENTS**

July 15 – 18, 2018

**Texas Pecan Growers Conference and Trade
Show, Embassy Suites and Conference Center
San Marcos, TX**

**Contact: TPGA at: 979-846-3285 or
pecans@tpga.org**

August 24, 2018

**Arizona Pecan Growers Annual Conference
Desert Diamond Casino and Hotel
Tucson, AZ**

**Contact: Mike Kilby: mkilby@email.arizona.edu
or 520-403-4613**

September 6, 2018

**Georgia Pecan Growers Assoc. Annual Fall Field
Day**

**USDA-ARS Facility, Byron, GA.
Contact GPGA @ 229-382-2187**

The information given herein is for educational purposes only. References to commercial products or trade names are made with the understanding that no endorsement by the Texas A&M AgriLife Extension Service is implied. The members of Texas A&M AgriLife will provide equal opportunities in programs and activities, education, and employment to all persons regardless of race, color, sex, religion, national origin, age, disability, genetic information, veteran status, sexual orientation or gender identity and will strive to achieve full and equal employment opportunity throughout Texas A&M AgriLife. The Texas A&M University System, U.S. Department of Agriculture, and the County Commissioners Courts of Texas Cooperating

.....

Table 5. Suggested insecticide to control yellow pecan aphids (blackmargined and yellow pecan aphids) and black aphids. This information is provided for educational purposes only. Read and follow label directions.

Insecticide			
Active ingredient	IRAC group	Brand name	Remarks
Imidachloprid*	4A	Admire® Pro	Do not graze livestock
		Advise® 2 FL, Max	
		Alias 2F	
		AmTide Imidacloprid 2F	
		Amtide Imidacloprid 4F	
		Couraze® 1.6F	
		Couraze® 2F	
		Couraze 4F	
		Lada 2F	
		Imida® E-AG 1.6 F	
		Imida® E-AG 2F	
		Impulse® 1.6 F	
		Macho® 2.0 FL	
		Malice® 75WSP	
		Mana® Alias 4F	
		Merit® 2F, 75WSP	
		Montana® 2F	
		Nuprid® 1.6F	
		Nuprid® 2F	
		Nuprid® 4F	
Pasada® 1.6F			
Prey® 1.6			
Provado® 1.6F			
Sherpa®			
Trimax Pro®			
Widow®			
Wrangler			
Clothianidin	4A	Belay Arena	Do not graze
Flonicamid	9C	Beleaf 50SG, Carbine 50WG	Do not graze livestock, 40 day PHI
Pymetrozine	9B	Fulfill	Grazing allowed
Thiamethoxam	4A	Centric 40WG Flagship 25WG	Centric can be applied to bearing trees. Flagship can only be applied to nonbearing trees only.
Dimethoate	1B	Dimethoate® 4E Dimate® 4EC, 4E Dimethoate® 4EC Dimethoate® 5lb	Do not graze livestock in treated orchards. Marginal control of yellow aphids has been observed

Table 7. Suggested insecticides to control pecan leaf scorch mite. This information is provided for educational purposes. Read and follow label directions.

Insecticide			
Active ingredient	IRAC group	Brand name	Remarks
Fenbutatin-oxide	12B	Vendex® 50 WP	Do not apply within 14 days of harvest.
Dicofol	unknown	Kelthane® MF	Do not apply within 7 days of harvest
Hexythiazox	10A	Onager®	For non-bearing orchards only. Do not graze treated orchards.
		Savey® 50 DF	
		Hexygon DF	
		Hexy 2E	
Bifenazate	unknown	Acramite® 50 SC	Do not graze treated orchards.
Spirodiclofen	23	Envidor® 2 SC	Grazing allowed
Cyflumetofen	25	Nealta	Grazing allowed
Fenazaquin	21	Magus	Non-bearing trees only
Fenpyroximate	21A	Fujimite 5EC	Grazing allowed
		Portal XLO	Grazing allowed