

rangeland grass may be used for used for grazing or cut for forage 0 days after application. Do not cut grass to be dried and harvested for hay until 7 days after the last application. Restricted use insecticide.

**Baythroid XL.** 12.07% cyfluthrin. Fall armyworm and grasshoppers. Pasture, rangeland, grass grown for hay and seed. Zero days to grazing or harvesting hay. Restricted use insecticide.

**Dimilin 2L.** 22% diflubenzuron. Fall armyworm and immature grasshoppers. Dimilin must be applied before armyworm larvae reach ½ inch or larger. Provides residual control for up to 2-3 weeks, as long as forage is not removed from the field. Label does not list a restriction on grazing.

**Prevathon.** 5% chlorantraniliprole. Fall armyworm and grasshoppers. Prevathon has a 0 day waiting period for harvest or grazing and is not a restricted use insecticide.

**Besiege.** 9.26% chlorantraniliprole and 4.63% lambda cyhalothrin. Fall armyworm and grasshoppers. Pasture and rangeland grass may be used for grazing or cut for forage 0 days after application. Do not cut grass to be dried and harvested for hay until 7 days after the last application. Restricted use insecticide.

**Sevin 4F, Sevin XLR, Sevin 80S, Generic Carbaryl.** Fall armyworm and grasshoppers. When applied to pastures, there is a 14 day waiting period before grazing or harvesting.

**Malathion\_57% and Malathion ULV.** Fall armyworm and grasshoppers. Zero days to harvest or grazing.

**Intrepid 2F.** Fall armyworm (not grasshoppers). Begin applications when first signs of armyworm feedings appear. Use higher rates for heavier infestations. Do not harvest hay within 7 days of application. No pre-harvest interval for forage. 0 days to grazing.

**Tracer.** Treat when armyworm eggs hatch or when larvae are small. Use higher rates for larger larvae. Do not graze until spray is dry. Do not harvest hay or fodder for 3 days after treatment. Do not allow cattle to graze until spray has dried.