



# Special Local Need

SLN No. TX-180005

## Duplosan™ Herbicide

EPA Reg. No. 228-742

For Post-harvest destruction / removal of all cotton including 2,4-D  
and Dicamba tolerant varieties

**FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF TEXAS**

This SLN label expires on 12-31-2023 and must not be used or distributed after this date

### DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN ANY MANNER INCONSISTENT WITH ITS LABELING. THIS LABELING MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF APPLICATION. READ THIS SLN LABELING AND THE ENTIRE LABEL AFFIXED TO THE PRODUCT CONTAINER BEFORE APPLYING. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THIS SLN LABELING AND THE LABEL AFFIXED TO THE PRODUCT CONTAINER.

<b>USE FOR:</b>
Post-harvest destruction / removal of all cotton including 2,4-D and Dicamba tolerant varieties.
<b>SPRAY MIXTURE PREPARATION:</b>
Unless otherwise specified in the use specific instructions in this table, add one-half the required amount of water to the spray tank, then add this product with agitation, then add any tank mix partners and finally, add the balance of the water with continued agitation. This material forms an emulsion in water. Emulsions tend to separate on standing. Provide continuous agitation to prevent separation and to form a uniform mixture.
<b>RESTRICTIONS &amp; LIMITATIONS:</b>
<b>DO NOT</b> plant any crop other than those listed on this label within 30 days to fields treated with Duplosan® Herbicide. <b>DO NOT</b> exceed 1.5 lbs ae / A / application (3 pints of Duplosan® Herbicide) <b>DO NOT</b> make more than 2 applications / year. <b>DO NOT</b> harvest cotton treated with Duplosan® Herbicide.
<b>POST HARVEST COTTON DESTRUCTION:</b>
Mix 32 – 48 fl. oz. (2 – 3 pints) of Duplosan® Herbicide (1 – 1.5 lbs dichloprop-p acid equivalent) per acre treated. For best results, apply Duplosan 2 weeks after final cotton harvest. Some leaf regrowth on the mowed or standing stalks will increase product uptake and performance. Applications of Duplosan may commence immediately after harvest to freshly shredded stalks or standing stalks, but this may result in less consistent control. Cotton stalks should not be shredded within 2 weeks after application due to reduced translocation to the root and reduced efficacy. A second application after the initial treatment may be required for complete kill. Add crop oil concentrate at 1% v/v.

### NOTICE

Read the "WARRANTY DISCLAIMER" and "LIMITATION OF LIABILITY" in the label booklet for this product before using this product. Those terms apply to this supplemental labeling and if those terms are not acceptable, return the product unopened at once.

Manufactured for Nufarm Americas Inc.  
11901 S. Austin Ave, Alsip, IL 60803